

## **Historic, archived document**

Do not assume content reflects current scientific knowledge, policies, or practices.

U. S. DEPARTMENT OF AGRICULTURE.

---

FARMERS' BULLETIN No. 206.

---


# MILK FEVER:

## ITS SIMPLE AND SUCCESSFUL TREATMENT.

BY

JOHN R. MOHLER, V. M. D.,

*Chief of Pathological Division, Bureau of Animal Industry.*



WASHINGTON:  
GOVERNMENT PRINTING OFFICE.

1904.

## LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,  
BUREAU OF ANIMAL INDUSTRY,  
*Washington, D. C., August 17, 1904.*

SIR: Under the date of May 18, 1904, Circular No. 45, entitled "Milk Fever: Its Simple and Successful Treatment," by Dr. John R. Mohler, Chief of the Pathological Division of this Bureau, was issued in order to acquaint the owners of cows with a new method of treatment for this disease which has proved to be very successful and at the same time quite simple. The demand for this circular has been such that it is deemed advisable to issue it as a Farmers' Bulletin in order that the information contained in it may have the still wider circulation which is afforded by this popular series of bulletins. I therefore recommend its publication as a Farmers' Bulletin.

Respectfully,

A. M. FARRINGTON,  
*Acting Chief of Bureau.*

Hon. JAMES WILSON,  
*Secretary of Agriculture.*

## CONTENTS.

	Page.
Preliminary remarks .....	3
Name and synonyms .....	3
Description of disease.....	3
Predisposition and cause.....	4
Symptoms .....	6
Appearance after death .....	8
Prognosis and mortality.....	8
Treatment .....	8
The potassium iodide treatment .....	9
The new air treatment.....	10
Prevention.....	14

## ILLUSTRATIONS.

	Page.
FIG. 1. Cow affected with milk fever.....	6
2. Apparatus for injecting sterile atmospheric air into the udder.....	12

# MILK FEVER: ITS SIMPLE AND SUCCESSFUL TREATMENT.

---

## PRELIMINARY REMARKS.

Milk fever is a very common, and until recently a frequently fatal, disease affecting cows in all the large dairy districts of this and other countries. As it usually attacks the best milking members of the herd and at a time when the milk flow is the heaviest, the malady is one which has caused very severe losses to our dairy industry. It is therefore of the greatest economic importance that every milk producer acquaint himself with the present extremely successful methods of treating this disease, especially the injection of filtered atmospheric air into the udder. This form of treatment has been adopted within a comparatively recent time, and, in view of the uniform success that has followed, every dairyman should become familiar with its use and should provide himself with a suitable apparatus for its application, especially if he is located where the services of a competent veterinarian can not be secured. This method of dealing with the disease does not make the assistance of the veterinarian undesirable, in case it is obtainable, as the professional man may frequently be of the greatest assistance in treating complicated symptoms, should they arise.

## NAME AND SYNONYMS.

The common name for this malady—milk fever—is an erroneous and misleading one, as in reality fever is usually absent; instead, there is generally an actual reduction in body temperature. A far better and more distinctive term and one that describes the actual condition much more precisely is parturient paresis. The disease has also several other names in various parts of the country, such as calving fever, parturition fever, parturient apoplexy, parturient collapse, puerperal fever, vitular fever, and dropping after calving.

## DESCRIPTION OF DISEASE.

Milk fever is a disease of well-nourished, plethoric, heavy-milking cows; it occurs during the most active period of life (fourth to sixth calf), and is characterized by its sudden onset, and the complete paralysis of the animal with loss of sensation, and by following closely the act

of calving, or parturition, terminating in a short time in recovery or death. One attack predisposes the animal to a recurrence of the trouble. While this disease may occur at any time during the whole year, it is seen principally during the warm summer season. The affection is almost entirely confined to the cow, although a few cases have been reported in the sow and goat. Sheep are entirely free from the disease.

### **PREDISPOSITION AND CAUSE.**

There are few diseases among our domesticated animals regarding the exact cause of which more widely different theories have been advanced than that of milk fever. The causes may properly be divided into two kinds—predisposing and direct. Experience shows one of the most prominent predisposing causes to be the great activity of the milk-secreting structure, namely, the udder. This organ is most active after the fourth, fifth, and sixth parturition, and this is the time of life when the vast majority of cases occur. The disease is almost unknown in heifers with the first calf and decreases in frequency steadily after the most active milking period is past. It is rarely, if ever, met with in pure beef breeds, such as the Shorthorn, Angus, and Hereford, while its main inroads are made into the heavy-milking breeds, such as the Holstein, Jersey, and Guernsey. Another factor that is probably of equal importance with the activity of the udder in producing the disease is the existence of a plethoric condition of the system, the result of excessive feeding and lack of exercise before calving. In heavy-milking cows all the food eaten in excess of that required to make up for the normal waste of the system is turned into milk and not used for the laying on of flesh or fat. Fleshiness is therefore an unnatural condition in these animals, and the period during which they are “dry” is usually very short; indeed, many of these cows continue to secrete milk right up to the time of calving. In those cases where the animals go dry the excess of nutriment in the food has no avenue of escape and immediately becomes stored up in the glands and in the blood, throwing the system into a high state of plethora. Now, at the time of calving all the blood which has been supplying the fetus is suddenly thrown back on the circulation, and if the udder does not begin active secretion very promptly plethora becomes extreme. The blood plasma under these conditions is very rich and dense, containing a large percentage of albumen and glycogen, and causing a shrinkage in size of the blood cells. This condition is invariably seen when the blood of milk-fever patients is examined under the microscope.

Fatness of the animal has been ascribed an important place among the causes of milk fever. This, however, in itself is probably not a predisposing cause. The beef breeds (Angus and Shorthorn) are

usually in far better condition at the time of calving than the milking breeds (Jersey and Holstein), and yet milk fever is a rarity in the former. At the same time it must be understood that a fat Jersey is more predisposed than one poor in flesh. In the fat Jersey the system is already loaded with an excess of nutriment, and, at the time of calving, extreme plethora is more readily produced than in the thin animal where the excess of nutritive elements could be more readily used and stored in the depleted muscular and glandular structures of the body. Fatness is therefore only of importance in the production of the disease in so far as it tends to increase glandular activity, particularly of the udder, and because of the higher state of plethora of the fat animal.

Regarding the direct cause of milk fever numerous theories have been advanced by various investigators, but only to be abandoned as further discoveries in pathology were made. Thus Schmidt, of Mühlheim, basing his theory upon the striking resemblance of the symptoms of milk fever to those of sausage poisoning, claimed that the former was due to an auto-intoxication, produced by the absorption of toxins from the uterus. This was a great advance over the theories which up to this time had been considered. Nevertheless, the medication recommended by this writer, which aimed at the antiseptic treatment of the womb, failed to decrease the great mortality of the disease, and his theory was finally entirely superseded in 1897 by that of J. Schmidt, of Kolding, Denmark. This able investigator first directed attention toward the udder by claiming (as Schmidt, of Mühlheim, had done) that the disease was an auto-intoxication, but produced by the absorption from the udder of leucomaines, resulting from the decomposition of the first milk (colostrum). Following up this idea Schmidt considered that the treatment should be directed toward retarding the secretion of the udder and at the same time neutralizing the leucomaines, or toxins, already present, by the use of some anti-toxic agent. He therefore advocated the injection of the udder with an aqueous solution of potassium iodide, which method was followed by an immediate decrease in the mortality to a very marked degree. The great success attendant upon this line of treatment at once gave the theory general recognition, and this very difficult problem was at last thought to be solved. Within the last few years the injections of etherized air, oxygen, and sterile atmospheric air have been used with wonderful success; reducing the death rate much lower than the potassium iodide had done. It therefore seems that a thorough distention of the udder is far more important than the antitoxic action of potassium iodide.

In explanation of the efficiency of this treatment some writers have claimed that after calving the determination of blood is suddenly

changed from the uterus to the udder, which produces an anemia (bloodlessness) of the brain. By thoroughly distending the udder the pressure therein causes a decrease in its circulation, which tends to equalize the distribution of the blood in other parts of the body, thus relieving this anemia of the brain and the consequent symptoms of milk fever. It has also been suggested that the highly favorable results obtained by the injection of atmospheric air and other gases into the udder are due to the fact that milk fever is produced by an anaerobic organism (a germ unable to live in the presence of air), which invades the udder. If this should prove true it is probable that this organism remains localized in the udder, as the bacillus of lockjaw remains localized at the point of injury, and produces a highly potent toxin, which, when absorbed into the system, has a specific action on the nerve centers. These theories, however, have not been proved, and the determination of the cause of this affection still requires investigation.




FIG. 1.—Cow affected with milk fever.

### SYMPTOMS.

This disease in its typical and most common form is comparatively easy to diagnose and one which almost every dairyman knows immediately before the arrival of the veterinarian. It usually comes on within two days after the birth of the calf and is practically never seen after the second week. In isolated instances it has been observed a few days before calving. At the commencement of the attack there is usually excitement; the cow is restless, treads with the hind feet, switches the tail, stares anxiously around the stall or walks about uneasily. She may bellow occasionally, show slight colicky symptoms, and make ineffectual

attempts at relieving the bowels. These symptoms are rarely recognized by the owner, but they are followed within a few hours by beginning paralysis, indicated by a staggering gait, especially in the hind legs, and by weakening of the knees and fetlocks in front. The patient now becomes quieter, the gait more staggering and weak, and finally the animal goes down and is unable to rise. The paralysis by this time is general, the calf is unnoticed, and the cow lies perfectly quiet with the eyes partly closed and staring and showing a complete absence of winking when the eyeball is touched. She is absolutely unheeding of her surroundings and flies may alight with impunity on all parts of the body without causing the slightest movement to dislodge them. While down the patient assumes a very characteristic position (see fig. 1), which is of great aid in diagnosis. The head is turned around to the side (usually the left) and rests on the chest, causing a peculiar arching of the neck. If the head is drawn out straight, it immediately flops around to the side again when the force is removed. The body usually rests slightly to one side, with the hind legs extended forward and outward and the fore legs doubled up in their normal position. There is paralysis of the muscles of the throat, so that swallowing is impossible, and in case drenching is attempted there is great danger of the fluids going into the lungs and setting up traumatic pneumonia. Paralysis of the rectum and bladder is also complete and the movement of the intestines is so suppressed that purgatives are frequently powerless to reestablish it. Fermentation in the paunch with consequent bloating is sometimes seen, particularly when the patient is allowed to be stretched out on her side. The secretion of milk is diminished and may be suspended entirely. Sugar is voided in the urine, depending in quantity on the severity of the attack. The pulse is weak and at times hardly perceptible to the finger, averaging from 50 to 70 beats per minute. Later in the disease, however, and especially in those cases with unfavorable terminations, it may reach 100 per minute. There is seldom noticed a rise of temperature. Sometimes at the commencement of the attack the temperature may reach 103° F., but there is a steady decrease to as low as 95° F. as the disease progresses. The temperature rapidly rises again as improvement is manifested. Convalescence occurs rapidly, and on the day following the onset of the disease, and in some cases even within a few hours, the animal may be up eating and drinking in a normal manner. Sometimes, however, a slight paralysis of the hind quarters persists, and may remain for a week or even longer, indicating that some structural change must have occurred in the nerve centers. In fatal cases the animal may remain perfectly quiet and die in a comatose condition from complete paralysis of the nervous system, but more frequently there is some agitation and excitement prior to death with tossing



about of the head. Death, like recovery, usually occurs in from eighteen to seventy-two hours after the onset of the malady.

### **APPEARANCE AFTER DEATH.**

The postmortem appearances in an animal dead of this disease are frequently entirely negative and not in the slightest degree characteristic. This further upholds the theory that milk fever is an intoxication and not a bacterial infection, as in the latter case the lesions would be more marked and distinguishable. The postmortem also fails to substantiate the fermentation theory of Schmidt-Mühlheim, as the uterus is generally found contracted and its mucous membrane intact. The third stomach is sometimes found impacted with dry, hard masses of food, and there may be some fermentation in the intestines. Sugar in varying percentage is always found in the urine in the bladder as well as in that drawn prior to death. Various particles of food may be found in the larynx together with congestion and swelling of the mucous membrane of the trachea and bronchi. Pneumonia, traumatic in origin, may also be observed if drenching has been attempted after paralysis of the throat has occurred. The blood usually appears dark and thick and congestion of the spinal cord and of the base of the brain is also quite frequently present.

### **PROGNOSIS AND MORTALITY.**

Prior to the introduction of the Schmidt treatment milk fever was considered an exceedingly serious malady and the death rate was placed by various authors at 40 to 50 and even 70 per cent. However, after the introduction and general application of potassium iodide injections into the udder, the mortality was reduced in Denmark and Germany to 17 per cent, in Switzerland to 22 per cent, and in Austria to 25 per cent, while in this country the statistics collated at the Iowa Experiment Station show 119 recoveries without complication out of 166 cases, a mortality of 28 per cent. Since the use of sterile atmospheric air for the injection of the udder, the death rate is even much lower than with the potassium iodide treatment, and in Denmark out of 914 patients, 884, or 96.7 per cent, recovered. In general, the nearer the attack follows the act of calving the more severe it proves and the graver the danger. The severity also greatly increases with each subsequent attack.

### **TREATMENT.**

In the administration of medicine by the mouth, and especially drenches, great care should be taken to prevent the fluids from getting into the larynx and from there into the lungs where they will set up traumatic pneumonia, which is almost invariably fatal. In case the

throat is not paralyzed the drench may prove of value and should be given slowly and immediately stopped at the first sign of uneasiness or coughing on the part of the animal. While the patient lies on the side she must raise the weight of her body at each inspiration, which is very exhausting, and hypostatic congestion of the dependent lung is greatly favored. Consequently it is of importance that the cow should be kept propped up on the breastbone by means of bags of chaff or straw placed against her side. In the way of medicinal treatment purgatives may be given in the first stage of the disease when the animal can swallow, with the precautions above mentioned. One pound of Epsom salts and 2 ounces of creolin dissolved in a pint of water will prove beneficial. The creolin is added for its antiseptic action to prevent fermentation in the paunch with the consequent danger of the eructation of foods and their subsequent passage into the windpipe. Ammonium carbonate in 2-dram doses dissolved in 2 ounces of water will be found to act equally as well as the creolin in this respect. Epsom salts is rather slow in its action, and an injection under the skin with a hypodermic syringe of  $1\frac{1}{2}$  to 2 grains of eserine sulphate, when obtainable, will be found quicker and more efficacious. The rectum should be emptied and injections of 1 to 2 gallons of warm water given to stimulate intestinal movements. However, the normal movement of the bowels, once lost, is exceedingly hard to reestablish, and sometimes all efforts in that direction fail. The urine should be drawn with a catheter or by pressure on the bladder with the hand in the rectum, as the bladder is paralyzed and unable to empty itself.

The feeble pulse and subnormal temperature call for the administration of stimulants. Injections under the skin of 1 dram of the following solution every three hours are probably the most efficacious: 80 grains of caffeine, 60 grains of sodium salicylate, and 4 drams of water. Similar injections of 1 grain of strychnia sulphate three times daily will also be found very beneficial, although numerous other drugs may be recommended, as spirits of camphor, veratrin, tincture of digitalis, alcohol, etc. In case the animal is very excitable the head should be restrained in such a manner as to prevent injury, and, in case the violence becomes excessive,  $1\frac{1}{2}$  ounces of chloral dissolved in a quart of water may be injected into the rectum, or 5 grains of morphine sulphate under the skin.

#### THE POTASSIUM IODIDE TREATMENT.

As previously stated Schmidt, of Kolding, advanced the theory in 1897 that the cause of milk fever was the absorption of leucomaines from the udder, and recommended that potassium iodide be injected to prevent the formation of the toxin and to neutralize that already existing. This was the most rational theory so far advanced and the treat-

ment proved to be beneficial, being followed by astonishingly good results. After this treatment was generally resorted to throughout Europe and America, the death rate fell from 40 per cent to 17 per cent. The apparatus required for the Schmidt treatment is exceedingly simple and consists of a piece of rubber tubing about 4 feet long, to one end of which is attached an ordinary milking tube which is inserted into the teat. At the other end a funnel is fitted, into which the solution is poured. Previous to the injection the udder should be thoroughly milked out and washed off with warm water and soap, followed by a 5 per cent solution of carbolic acid or creolin. A clean towel should be placed under the udder to keep it from coming in contact with the stable litter or other filth. Two and one-half drams of potassium iodide are then added to one quart of water previously boiled for fifteen minutes and allowed to cool to the temperature of the body. The funnel and tubing should likewise be disinfected before the injection. The milking tube is inserted into the four teats in succession, each quarter of the udder, after it has been milked out clean, receiving one-half pint of the liquid. The udder should then be thoroughly massaged to make sure that all the milk canals are penetrated by the liquid. In case improvement does not occur the injection may be repeated once or twice at intervals of eight hours, always observing the same antiseptic precautions, as it is possible to produce a dangerous mammitis (caked bag) and ruin the udder by careless injections which introduce pathogenic bacteria. This danger, however, is entirely obviated by the use of ordinary antiseptic precautions, as described above.

#### THE NEW AIR TREATMENT.

Of all known methods of treating milk fever, the injection of sterile atmospheric air into the udder is by far the most simple and practicable as well as the most efficacious and harmless one at our disposal, and only occasionally requires that medicinal treatment be given.


For a considerable length of time the entire value of Schmidt's treatment was considered to be the antitoxic action of potassium iodide, and soon numerous investigators began injecting various other antiseptics, such as carbolic acid, creolin, etc., with equally good results. Sterile water and sterile salt solution were tried with no increase in the mortality, and it was therefore considered that the distention of the udder was as important a factor as the antitoxic action of the iodide of potash. Continuing along these lines, Kortman used antiseptic gases (etherized air) with beneficial results. Oxygen was then tried by Knüsel with increasing success and the deaths among the experimental cases virtually ceased. The apparatus for treating with oxygen and etherized air, however, are expensive and cumbersome, and this greatly limits their use by the average practitioner.

To Andersen, of Skanderborg, belongs the credit of first having made use of plain atmospheric air, although Schmidt had previously recommended the admittance of air with the potassium iodide solution for the purpose of obtaining greater diffusion of the liquid. Andersen first injected air along with sterile water and then by itself. The results were astonishingly successful. Thus Schmidt reports that out of 914 cases treated in Denmark, 884, or 96.7 per cent, were restored to health. The record of 140 of these animals shows that recovery occurred in the average time of  $6\frac{2}{3}$  hours. Of this number 25 cases required a second injection, while in 3 of the latter number it was necessary to give a third treatment before they were able to get upon their feet. The treatment is also practically harmless, as the statistics of the above-mentioned 914 patients show that only 1 cow was affected with a severe attack of caked bag after this treatment, while in 4 other cows a milder inflammation of the udder was apparent. Equally good results have likewise been obtained in this country.

The method of injecting filtered air into the udder is easy of manipulation, requires but little time, and is readily accomplished by means of a milk-fever apparatus, such as is illustrated in fig. 2.<sup>a</sup> It consists of a metal cylinder *f* with milled screw-caps *c* and *d* on either end. Cap *c* may be removed in order to place sterile absorbent cotton within the chamber. To this cap the rubber bellows *a* and *b* are connected by 9 inches of rubber tubing. Cap *d* is to be removed together with the attached 18 inches of rubber hose, at the free end of which is the self-retaining milking tube *g*, for the purpose of disinfection before treating each case. The pulling on or off of the tubing on the nozzles of the milled caps is thus rendered unnecessary. Within the metal cylinder at *e* is a wire net, which prevents the obstruction of the outlet of the chamber by holding back the sterile cotton, and also permits of the unscrewing of the lower cap and the disinfection of this portion of the apparatus, including the milking tube, without contaminating the packing. Absorbent cotton impregnated with carbolic acid (carbolyzed cotton) or other suitable disinfectant can be purchased from the drug trade in most localities, and is better, though slightly more expensive, than the plain cotton.

Previous to making the air injection, the hands of the operator should be thoroughly cleansed and the udder should receive the same careful antiseptic treatment as has been recommended in discussing the injection of potassium iodide. Soap and water should be applied to the teats and udder, after which they should be carefully disinfected with a 5 per cent solution of carbolic acid (3 tablespoonfuls of pure

<sup>a</sup> Persons desiring the name and address of firms manufacturing milk-fever outfits, together with the price of same, will be furnished with such information upon application to the Chief of the Bureau of Animal Industry.



carbolic acid to 1 quart of water). A clean towel should then be placed under the udder to prevent the teats from coming in contact with dirt or filth of any kind. The milking tube, before it is placed in the teat, should have been perfectly sterilized by boiling for fifteen minutes, with the lower hose and cap of the cylinder attached, and the apparatus should be wrapped in a clean towel, without touching the milking tube, to prevent contamination before use. If the apparatus has been subjected to this treatment shortly before and it is desired to disinfect only the milking tube, the latter may be placed in a 5 per cent solution of carbolic acid for five minutes. It is then carefully inserted into the milk duct of the teat without emptying the udder of milk. Air is now pumped from the bulb *a* into the reservoir *b*, and thus a continuous flow of air is forced through the filtering chamber and into the udder. Slight massage or kneading of the udder will cause the innermost recesses of the milk tubules to become distended with the injected air. After one-quarter of the udder is well distended the milking tube is removed, care being taken to prevent the outflow of air by having an assistant tie a broad piece of tape about the teat at the time the milking tube is withdrawn. The same treatment is repeated with the other three teats until the udder is satisfactorily distended. In case the air becomes absorbed and no improvement is noted within five hours, a repetition of this treatment should be made under the same antiseptic precautions as at first. The tape should be removed from the teats two or three hours after the cow gets on her feet, the constricting muscles at the tip of the teats being now depended on for retaining the air. In this manner the air may be left in the udder for twenty-four hours, and when recovery is assured, it should be gradually milked out. It is needless to say that the calf should not be permitted to suck during this period.

Inflammation of the udder (caked bag) is avoided if the milking tube is thoroughly disinfected before each application, and if the cow's teats and bag and the hands of the operator have been properly cleansed. If the apparatus is kept in its case free from dust and dirt, the absorbent or medicated cotton in the metal cylinder will efficiently filter enough air to distend the udders of six cows. After this number has been treated it is advisable to replace the old cotton with a fresh sterile supply, which should be placed loosely in the cylinder.<sup>a</sup>

While this method of treating milk fever is a comparatively easy one for a farmer or dairyman to adopt, he can not expect to have the

---

<sup>a</sup>A number of bacteriological tests were made in order to determine the efficiency of the absorbent cotton in the cylinder to filter out the microorganisms from the atmospheric air. For this purpose agar plates and slant agar tubes were used and the results obtained proved the absolute certainty of this quantity of cotton filtering successfully a sufficient volume of air to distend six good-sized udders.

same successful results as those obtained by a skilled veterinarian, and it is therefore advisable that the services of such a veterinarian should always be obtained in those districts where it is possible. In many cases it will be found that the injection of air into the udder will be sufficient to combat the disease without any other treatment, but it is always advisable to study the symptoms of each individual case and administer in a rational manner the indicated medicines.

### PREVENTION.

Until recently most stringent measures were resorted to by every careful dairyman to prevent the development of the disease in his herd. However, since the treatment of the present day has so greatly reduced, and even in some cases obliterated, the mortality, prevention is no longer such an important problem and therefore preventive measures which have a severe and lasting effect upon the animals should be abandoned from an economic standpoint. It has long been advocated to starve all suspected animals for two weeks prior to the birth of the calf. It is frequently noted that this has an injurious effect on the milk flow of the animal, from which it may require several weeks for her to recover and gain her normal output of milk. This measure is no longer considered advisable, as it is better to have cows attacked with the disease once in a while (the mortality being less than 5 per cent) than to decrease the flow from every heavy-milking cow for one to three weeks after she comes fresh by starving her before calving.

A method which is not quite so sure of reducing the plethoric condition of the cow, but which nevertheless proves very efficient and is without the slightest permanent injurious effect, is the administration of 1 to 1½ pounds of Epsom salts two or three days prior to calving. In case this has been neglected and a well-nourished, heavy-milking cow has passed through an easy nonexhausting calf birth, the administration of the salts after the labor is over should by no means be neglected. Bloodletting has also been advocated, but there is always the danger of exciting the blood-making organs to excessive activity, thus largely neutralizing the effect. It should therefore be resorted to only when the cow is extremely fat, is a heavy milker, and has had one or more previous attacks. The blood should be drawn from the jugular vein until the pulse softens perceptibly, 1½ pints for every 100 pounds of the animal's body weight being about the right amount.

Another very good preventive measure, and one easily carried out, though frequently overlooked, is to give the cow plenty of exercise up to the time of calving. Many animals are allowed to run continuously on pastures from the time they go dry until a week or two before calving, when they are transferred to the stable without any subsequent

exercise. This is very conducive to the enriching of the blood and the development of the disease.

The most recent preventive treatment suggested is in line with the favorable results obtained by the injection of air into the udder. It consists in allowing the susceptible cow to retain in the udder for 24 hours after calving all the milk except the small quantity required by the calf, which should be taken if possible from each quarter. The distention of the udder naturally follows as in the air treatment and acts as a preventive against milk fever. In the Island of Jersey and at the Biltmore Farms, N. C., where this practice is common, the number of milk fever cases has been greatly lessened. General sanitary conditions should also be looked after, such as the supply of pure air and clean stabling, with plenty of clear cool water and laxative foods, such as grasses and roots. Some observers who believe in the microbic origin of the disease have recommended the cleaning of the manure and dirt from the animal and spraying the hind quarters and genitals with a 4 to 5 per cent solution of carbolic acid, lysol, or creolin, just prior to calving. From our present knowledge of the disease, however, this is probably unnecessary.



## FARMERS' BULLETINS.

The following is a list of the Farmers' Bulletins available for distribution, showing the number and title of each. Copies will be sent free to any address in the United States on application to a Senator, Representative, or Delegate in Congress, or to the Secretary of Agriculture, Washington, D. C. Numbers omitted have been discontinued, being superseded by later bulletins.

No. 16. Leguminous Plants. No. 22. The Feeding of Farm Animals. No. 24. Hog Cholera and Swine Plague. No. 25. Peanuts: Culture and Uses. No. 27. Flax for Seed and Fiber. No. 28. Weeds: And How to Kill Them. No. 29. Souring and Other Changes in Milk. No. 30. Grape Diseases on the Pacific Coast. No. 31. Alfalfa, or Lucern. No. 32. Silos and Silage. No. 33. Peach Growing for Market. No. 34. Meats: Composition and Cooking. No. 35. Potato Culture. No. 36. Cotton Seed and Its Products. No. 37. Kafir Corn: Culture and Uses. No. 38. Spraying for Fruit Diseases. No. 39. Onion Culture. No. 41. Fowls: Care and Feeding. No. 42. Facts About Milk. No. 43. Sewage Disposal on the Farm. No. 44. Commercial Fertilizers. No. 45. Insects Injurious to Stored Grain. No. 46. Irrigation in Humid Climates. No. 47. Insects Affecting the Cotton Plant. No. 48. The Manuring of Cotton. No. 49. Sheep Feeding. No. 50. Sorghum as a Forage Crop. No. 51. Standard Varieties of Chickens. No. 52. The Sugar Beet. No. 54. Some Common Birds. No. 55. The Dairy Herd. No. 56. Experiment Station Work—I. No. 57. Butter Making on the Farm. No. 58. The Soy Bean as a Forage Crop. No. 59. Bee Keeping. No. 60. Methods of Curing Tobacco. No. 61. Asparagus Culture. No. 62. Marketing Farm Produce. No. 63. Care of Milk on the Farm. No. 64. Ducks and Geese. No. 65. Experiment Station Work—II. No. 66. Meadows and Pastures. No. 68. The Black Rot of the Cabbage. No. 69. Experiment Station Work—III. No. 70. Insect Enemies of the Grape. No. 71. Essentials in Beef Production. No. 72. Cattle Ranges of the Southwest. No. 73. Experiment Station Work—IV. No. 74. Milk as Food. No. 75. The Grain Smuts. No. 77. The Liming of Soils. No. 78. Experiment Station Work—V. No. 79. Experiment Station Work—VI. No. 80. The Peach Twig-borer. No. 81. Corn Culture in the South. No. 82. The Culture of Tobacco. No. 83. Tobacco Soils. No. 84. Experiment Station Work—VII. No. 85. Fish as Food. No. 86. Thirty Poisonous Plants. No. 87. Experiment Station Work—VIII. No. 88. Alkali Lands. No. 89. Cowpeas. No. 91. Potato Diseases and Treatment. No. 92. Experiment Station Work—IX. No. 93. Sugar as Food. No. 94. The Vegetable Garden. No. 95. Good Roads for Farmers. No. 96. Raising Sheep for Mutton. No. 97. Experiment Station Work—X. No. 98. Suggestions to Southern Farmers. No. 99. Insect Enemies of Shade Trees. No. 100. Hog Raising in the South. No. 101. Millets. No. 102. Southern Forage Plants. No. 103. Experiment Station Work—XI. No. 104. Notes on Frost. No. 105. Experiment Station Work—XII. No. 106. Breeds of Dairy Cattle. No. 107. Experiment Station Work—XIII. No. 108. Saltbushes. No. 109. Farmers' Reading Courses. No. 110. Rice Culture in the United States. No. 111. Farmers' Interest in Good Seed. No. 112. Bread and Bread Making. No. 113. The Apple and How to Grow It. No. 114. Experiment Station Work—XIV. No. 115. Hop Culture in California. No. 116. Irrigation in Fruit Growing. No. 118. Grape Growing in the South. No. 119. Experiment Station Work—XV. No. 120. Insects Affecting Tobacco. No. 121. Beans, Peas and other Legumes as Food. No. 122. Experiment Station Work—XVI. No. 123. Red Clover Seed: Information for Purchasers. No. 124. Experiment Station Work—XVII. No. 125. Protection of Food Products from Injurious Temperatures. No. 126. Practical Suggestions for Farm Buildings. No. 127. Important Insecticides. No. 128. Eggs and their Uses as Food. No. 129. Sweet Potatoes. No. 131. Household Tests for Detection of Oleomargarine and Renovated Butter. No. 132. Insect Enemies of Growing Wheat. No. 133. Experiment Station Work—XVIII. No. 134. Tree Planting in Rural School Grounds. No. 135. Sorghum Sirup Manufacture. No. 136. Earth Roads. No. 137. The Angora Goat. No. 138. Irrigation in Field and Garden. No. 139. Emmer: A Grain for the Semiarid Regions. No. 140. Pineapple Growing. No. 141. Poultry Raising on the Farm. No. 142. Principles of Nutrition and Nutritive Value of Food. No. 143. The Conformation of Beef and Dairy Cattle. No. 144. Experiment Station Work—XIX. No. 145. Carbon Bisulphid as an Insecticide. No. 146. Insecticides and Fungicides. No. 147. Winter Forage Crops for the South. No. 148. Celery Culture. No. 149. Experiment Station Work—XX. No. 150. Clearing New Land. No. 151. Dairying in the South. No. 152. Scabies in Cattle. No. 153. Orchard Enemies in the Pacific Northwest. No. 154. The Home Fruit Garden: Preparation and Care. No. 155. How Insects Affect Health in Rural Districts. No. 156. The Home Vineyard. No. 157. The Propagation of Plants. No. 158. How to Build Small Irrigation Ditches. No. 159. Scab in Sheep. No. 161. Practical Suggestions for Fruit Growers. No. 162. Experiment Station Work—XXI. No. 164. Rape as a Forage Crop. No. 165. Culture of the Silkworm. No. 166. Cheese Making on the Farm. No. 167. Cassava. No. 168. Pearl Millet. No. 169. Experiment Station Work—XXII. No. 170. Principles of Horse Feeding. No. 171. The Control of the Codling Moth. No. 172. Scale Insects and Mites on Citrus Trees. No. 173. Primer of Forestry. No. 174. Broom Corn. No. 175. Home Manufacture and Use of Unfermented Grape Juice. No. 176. Cranberry Culture. No. 177. Squab Raising. No. 178. Insects Injurious in Cranberry Culture. No. 179. Horseshoeing. No. 181. Pruning. No. 182. Poultry as Food. No. 183. Meat on the Farm—Butchering, Curing, etc. No. 184. Marketing Live Stock. No. 185. Beautifying the Home Grounds. No. 186. Experiment Station Work—XXIII. No. 187. Drainage of Farm Lands. No. 188. Weeds Used in Medicine. No. 189. Information Concerning the Mexican Cotton Boll Weevil. No. 190. Experiment Station Work—XXIV. No. 191. The Cotton Bollworm. No. 192. Barnyard Manure. No. 193. Experiment Station Work—XXV. No. 194. Alfalfa Seed. No. 195. Annual Flowering Plants. No. 196. Usefulness of the American Toad. No. 197. Importation of Game Birds and Eggs for Propagation. No. 198. Strawberries. No. 199. Corn Growing. No. 200. Turkeys. No. 201. Cream Separator on Western Farms. No. 202. Experiment Station Work—XXVI. No. 203. Canned Fruits, Preserves, and Jellies. No. 204. The Cultivation of Mushrooms. No. 205. Pig Management. No. 206. Milk Fever. No. 207. Game Laws, 1904. No. 208. Varieties of Fruits Recommended for Planting. No. 209. Controlling the Boll Weevil in Cotton Seed and at Ginneries.